

*A periodic table will also be provided for the exam

$$\pi = MRT$$

$$\Delta T = k_f m$$

$$\Delta T = k_b m$$

$$R = 0.08206 \frac{L \cdot atm}{mol \cdot K}$$

$$P = X_{solvent} P^{\circ}_{solvent}$$

$$[A]_t = -kt + [A]_o$$

$$t \frac{1}{2} = \frac{[A]_o}{2k}$$

$$\ln[A]_t = -kt + \ln[A]_o$$

$$t \frac{1}{2} = \frac{0.693}{k}$$

$$\frac{1}{[A]_t} = kt + \frac{1}{[A]_o}$$

$$t \frac{1}{2} = \frac{1}{k[A]_o}$$

$$S_{gas} = k_H P_{gas}$$

$$1 \text{ atm} = 760 \text{ mm Hg} = 760 \text{ torr}$$

$$k = Ae^{-E_a/RT}$$

$$\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$R = 8.314 \frac{J}{mol \cdot K}$$

$$K = ^{\circ}C + 273$$

1. Rank the following solutions in terms of increasing boiling point (lowest to highest boiling point)

0.100 m CaCl_2 , 0.40 m $\text{C}_6\text{H}_{12}\text{O}_6$, 0.125 AlCl_3 , 0.2 m $\text{C}_{12}\text{H}_{22}\text{O}_{11}$

Order (low to high bp)

2. Concentrated HCl is 37% by mass HCl with a solution density of 1.2 g/mL.

a. What is the molality of the solution?

b. What is the molarity of the solution?

3. Myoglobin is a protein that uses a heme group to bind oxygen. A sample of myoglobin (a non-electrolyte) with a mass of 1.0 grams is dissolved in water for a final volume of solution of 100.0 mL. The osmotic pressure of the solution is 11.0 torr at 25.0°C. Find the molar mass of myoglobin.


4. A certain reaction has a rate constant of $1.75 \times 10^{-1} \text{ s}^{-1}$ at 20.0°C . What is the value of the rate constant (k) at 40.0°C if $E_a = 55.5 \text{ kJ/mol}$?

5. A student wants to study the decomposition of a gas "A", and obtained the following data in the lab.

Time (s)	[A], (M)
0	1.00×10^{-2}
60	6.83×10^{-3}
120	5.18×10^{-3}
180	4.18×10^{-3}
240	3.50×10^{-3}
300	3.01×10^{-3}
360	2.64×10^{-3}

- a. What is the order of the reaction with respect to A? BRIEFLY explain your reasoning.
- b. What is the differential rate law including the value of the rate constant with units?
- c. What is the concentration of [A] after 500 seconds if the initial concentration $[A]_0$ was 0.0100 M?

6. Given the following data below for the reaction with the rate defined as $\Delta[\text{I}^-]/\Delta t$



	1	2	3
[I ⁻]	0.10	0.20	0.30
[OCl ⁻]	0.050	0.050	0.010
Rate (M/ s ⁻¹)	3.05×10^{-4}	6.20×10^{-4}	1.83×10^{-4}

a. Determine the rate law for the reaction



b. What is the rate constant with units for the reaction?

7. You study the following reaction in the lab: $\text{NO}_2(g) + \text{CO}(g) \rightarrow \text{NO}(g) + \text{CO}_2(g)$

The experimental rate law is $\text{rate} = k[\text{NO}_2]^2$

Answer the following questions regarding this reaction.

a. Is the following reaction mechanism consistent with the experimentally observed rate law, $\text{rate} = k[\text{NO}_2]^2$? Explain.




b. Please write a rate law for each elementary step in part a.

c. Circle the intermediate in the mechanism in part a

d. Does this reaction mechanism use a catalyst? (Circle one) YES OR NO


8. Please label the following reaction coordinate diagram below

Reactants (R), products (P), transition state 1 (TS1), transition state 2 (TS2), activation energy for step 1 (Ea1), activation energy for step 2 (Ea2), intermediate (I), and ΔH (including whether exo- or endothermic)




9.

For the reaction



If you begin the reaction with 1.0 mol of N_2 , 2.0 mol of H_2 , and sufficient C(s) in a 2.00 L container, what are the concentrations of N_2 , CH_3NH_2 , and H_2 at equilibrium?

10. For the equilibrium shown below, K_p at 1100°C for this process is 1.13×10^{-5} . If a sample with an initial Cl_2 gas pressure of 0.500 atm was allowed to reach equilibrium. What is the partial pressure of each gas at equilibrium AND the total pressure in the reaction vessel?



11. A certain species of fish requires a dissolved oxygen concentration of $7.5 \text{ mg O}_2 / \text{L}$ of water. Suppose the temperature of a lake in summer is 28.0°C . Would the fish be able to survive in this lake? The Henry's law constant (k_H is $1.2 \times 10^{-3} \text{ M/atm}$) at this temperature. The partial pressure of oxygen gas is 0.17 atm at 28.0°C . Please show all your work.

12. What is the vapor pressure if 50.0 grams of calcium chloride is dissolved in 500.0 mL of water (density of water is 1.0 g/mL) at 25.0 °C. The vapor pressure of water at 25.0 C is 23.8 torr.