

How grandmother effects plus individual variation in frailty shape fertility and mortality: Guidance from human–chimpanzee comparisons

Kristen Hawkes¹

Department of Anthropology, University of Utah, Salt Lake City, UT 84112

In the first paper to present formal theory explaining that senescence is a consequence of natural selection, W. D. Hamilton concluded that human postmenopausal longevity results from the contributions of ancestral grandmothers to the reproduction of their relatives. A grandmother hypothesis, subsequently elaborated with additional lines of evidence, helps explain both exceptional longevity and additional features of life history that distinguish humans from the other great apes. However, some of the variation observed in aging rates seems inconsistent with the tradeoffs between current and future reproduction identified by theory. In humans and chimpanzees, our nearest living relatives, individuals who bear offspring at faster rates do not cease bearing sooner. They continue to be fertile longer instead. Furthermore, within both species, groups with lower overall mortality rates have faster rates of increase in death risk with advancing age. These apparent contradictions to the expected life history tradeoffs likely result from heterogeneity in frailty among individuals. Whereas robust and frail alike must allocate investments between current and future reproduction, the more robust can afford more of both. This heterogeneity, combined with evolutionary tradeoffs and the key role of ancestral grandmothers they identify, helps explain aspects of human aging that increasingly concern us all.

evolution of senescence | heterogeneity of frailty | human life history | menopause | human aging

Long postmenopausal survival is a characteristic of our species. The use of life expectancy to compare human populations can obscure this fact because high infant and juvenile mortality kept all national life expectancies below 50 until the 20th century (1). As historical demography shows, girls that survived childhood usually lived long past menopause in previous centuries (2). Hunter-gatherer survival curves are especially instructive (3–7). They document characteristic human longevity in the absence of agriculture, public health institutions, and scientific medicine, all of which emerged long after the initial evolution of our species (8, 9). Distinctive and at first puzzling human postmenopausal survival was addressed in classic papers that used evolutionary theory to explain why living things grow old.

G. C. Williams (10) laid out demographic reasons why declines in adaptive performance with increasing adult age emerge from the forces of natural selection. Because life is risky, cohorts inevitably diminish across adulthood. Consequently, the forces of selection weaken with age as fewer remain to be affected by it at older ages. Williams explained how the same forces result in different rates of senescence among species that reproduce more than once depending on two aspects of life history. First, when background mortality risk is lower, more individuals survive to older ages and selection against senescence is stronger. Second, selection against senescence is also stronger when the potential fitness-related payoffs to survivors increase with age. He illustrated the latter effect with the slow senescence of indeterminate growers that continue to increase in size and rate of egg production throughout adulthood.

Concluding that evolutionary life history theory predicts no postreproductive period in normal lifespans, Williams then addressed the apparent contradiction posed by survival past meno-

pause in our own species by observing that older women still investing in descendants are not literally postreproductive. Hamilton (11) mathematically modeled the tradeoffs nominated by Williams and demonstrated that the forces of selection shape mortality schedules to converge asymptotically with the age when reproduction ends. This process leaves, as Williams had surmised, few if any postreproductives. Because “much the best” (ref. 11, p. 27) demographic data are available on humans, Hamilton used a human population to explore the fit of observation with theory. This required him to explicitly confront the apparent discrepancy in the case of humans (ref. 11, p. 37):

“... the rather definite age of menopause seems conspicuously ignored by the as yet gently rising curve of the force of mortality. It is, moreover, a matter of common knowledge that the post menopausal woman normally remains a useful and healthy member of the community for some time... [This] can be attributed to the beneficial effects of continued survival on the survival and reproduction of descendants... In fact... the comparatively healthy life of the postreproductive woman... inevitably suggests a special value of the old woman as a mother or grandmother during a long ancestral period...”

Such a grandmother hypothesis, subsequently elaborated with comparative and phylogenetic evidence not available when the classic papers appeared, can explain not only the evolution of human longevity but other similarities and differences in life history between humans and the other great apes. We live longer; we take longer to mature but have shorter birth intervals; and we share common ages of terminal female fertility with the other great apes (12, 13). The hypothesis focuses on females because as noted by both Williams and Hamilton our mid-life menopause is a central clue to human life history evolution and because the hypothesis employs E. L. Charnov’s (14, 15) model of tradeoffs faced by females to explain mammalian life history variation. The forces of selection explored by Williams (10, 16), Hamilton (11), Charnov (15), and many other students of life history evolution (17, 18) attend to fitness effects and not to proximate mechanisms, but T. B. L. Kirkwood’s disposable soma model (19) based on the same evolutionary tradeoffs between current and future reproduction has directed attention to processes of cellular maintenance and repair that affect somatic aging rates (20, 21). Such processes likely have similar effects in both sexes, because longer-lived mothers pass on their cellular maintenance mechanisms to both sons and daughters.

I briefly summarize this elaborated grandmother hypothesis, then turn to patterns that initially seem inconsistent with the tradeoffs between current and future reproduction identified in evolutionary

This paper results from the Arthur M. Sackler Colloquium of the National Academy of Sciences, “In the Light of Evolution IV: The Human Condition,” held December 10–12, 2009, at the Arnold and Mabel Beckman Center of the National Academies of Sciences and Engineering in Irvine, CA. The complete program and audio files of most presentations are available on the NAS Web site at www.nasonline.org/SACKLER_Human_Condition.

Author contributions: K.H. designed research, performed research, analyzed data, and wrote the paper.

The author declares no conflict of interest.

This article is a PNAS Direct Submission.

¹E-mail: hawkes@anthro.utah.edu.

explanations for senescence. I focus on two apparent inconsistencies between theoretical expectations and empirical observations. First, theory predicts that current reproductive output should subtract from effort invested in maintenance for survival and reproduction in the future, yet individuals with higher fertility rates tend to continue bearing offspring to older ages; and in humans, women with later last births then survive longer afterward (22–25). Second, theory predicts that lower adult mortality should slow rates of senescence, yet when populations of the same species are compared, the groups with lower mortality have steeper increases in death risk with advancing age (26, 27). More survival to older ages makes senescence—measured as the pace of increase in age-specific mortality—appear to be faster. Heterogeneity of frailty within populations may explain these apparent contradictions (28).

J. W. Vaupel and colleagues (29, 30) proposed that heterogeneity in frailty might explain why the increase in mortality rates across adulthood begins to slow and even cease at advanced ages in humans and many other taxa. If individuals vary in their vulnerabilities to death, the more frail will usually die younger. Survivors to the oldest ages will therefore be a subset of the population enriched with individuals that had lower vulnerability all along. L. D. Mueller, M. R. Rose, C. L. Rauser, and colleagues (31–33) judged Vaupel's hypothesis to be in conflict with Hamilton's forces and found those forces themselves sufficient to explain the mortality plateaus. I argue here that rather than being mutually exclusive alternatives, heterogeneity of frailty and tradeoffs between current and future reproduction explain different things. Both are needed to account for salient aspects of fertility and mortality schedules in general, and those of humans and chimpanzees in particular. As Williams and Hamilton recognized, women usually outlive their fertility. This is not true of chimpanzees. Although childbearing ends at the same age in both species, only humans regularly survive for decades longer. Heterogeneity within populations can explain why this divergence in life history results in fertility schedules with different shapes.

A Grandmother Hypothesis

Anthropologists continue to debate the phylogenetic relationships among fossil taxa representing our ancestors and cousins (34), but genetic evidence unequivocally corroborates Darwin's hypothesis about our African ape ancestry (35). The genera ancestral to our own are often characterized as bipedal apes (36), and chimpanzees are commonly used as a living model for the ancestors of our genus because they are genetically closest to us and similar in body and brain size to these extinct taxa (37). Correlations between life history traits and adult size across the living primates (15) support the relevance of a chimpanzee model for the early members of our lineage.

Like other primates, chimpanzees feed themselves after weaning (38). Systematic observations among modern hunter-gatherers show that human youngsters can be remarkably efficient foragers, acquiring large fractions of their own requirements at young ages (39–41); but unlike chimpanzees, humans still depend on provisioning by others after weaning. Help is especially crucial for certain kinds of foods (42). Reliance on resources that young juveniles cannot handle effectively requires mothers to provision weaned offspring, but mothers nursing new infants provide less for their weaned children who receive subsidies from grandmothers (43).

The productivity of Hadza hunter-gatherer grandmothers especially in gathering hard-to-acquire staples, and the importance of their subsidies to weaned children with infant siblings (43), suggests a scenario about the ancestral past. An ecological change that reduced the availability of foods juveniles could handle independently would have opened a novel fitness window to older females without nursing infants of their own (43). By helping to feed weanling grandchildren, elder females would have allowed their daughters to bear the next baby sooner without affecting the survival of previous offspring. More vigorous elders, through greater reproductive success of their daughters, would have spread

their slower somatic aging to more descendants. Longer adult lifespans then reduced the cost of waiting longer to mature, delaying age at maturity and increasing adult body size (44). Because later births would interfere with grandmothering, selection would not have favored delaying ages of fertility decline. Increased allocation to somatic maintenance would have left less for current reproduction through the childbearing years, but subsidies from elders would have more than compensated, raising the fertility of childbearers (45).

We hypothesized that such a shift might have given rise to genus *Homo* (45, 46) when drying environments and increased seasonality altered foraging opportunities for ancestral populations between 2 and 3 million years ago as forests shrank and grasslands spread across Africa (47, 48). Changes in body size and form are consistent with such a shift, as is the colonization of new habitats about that time. The hypothesis also helps explain the location of early archaeological sites and the composition of the faunal assemblages associated with them (49).

A formal model of the verbal grandmother scenario outlined here remains to be developed, but others have formalized links between the evolution of human longevity and the economic productivity of elders. H. S. Kaplan and A. J. Robson (50) have shown that aging rates can be connected to the contribution adults make to juvenile survival. R. D. Lee (51) has demonstrated that when intergenerational transfers of assistance are incorporated into a formal theory of senescence, it is the transfers instead of fertility that determine equilibrium aging rates. His simulations show that when elders transfer resources to close kin, mortality schedules very like those observed in hunter-gatherers are maintained by selection against deleterious mutations (52).

Age Structures

Our grandmother hypothesis relies on Charnov's model of life history evolution (14, 15) to explain how correlated allometries in mammalian life history features apply to humans (12, 53). Comparisons between other great apes and humans (13) have been essential in highlighting distinctive human life history features. As noted, chimpanzees are an especially important comparative model for phylogenetic, ecological, and morphological reasons. Fig. 1 shows the female side of the age structure for a human hunter-gatherer population and wild chimpanzees modeled from life tables.

The human example on the right in Fig. 1, the Hadza (6), is similar to other hunter-gatherers. Life expectancy at birth is <40 years, but a substantial fraction of adults are past the childbearing years. This is not true of chimpanzees, modeled on the left of Fig. 1 from the wild population synthesized from five wild study sites (54). Lower mortality in humans as compared to the other great apes has long been attributed to our propensity for cooperation and resource sharing (55), patterns that must surely affect death risks. The grandmother hypothesis highlights sharing by grandmothers in particular because, as noted by Hamilton, evidence that women remain healthy and productive past their fertility provides a clear link between human longevity and fitness payoffs to ancestral grandmothering. Sometimes elders survive with help from younger kin, but an evolutionary perspective predicts help to generally flow from older to younger relatives (56). Measures of strength and productivity among postmenopausal hunter-gatherers demonstrate their provisioning capacities (57, 58). High fractions of maximum function through and beyond the childbearing years in humans contrast with the earlier geriatric declines of chimpanzees (38, 59).

Demographic Aging Rates Between and Within Species

As expected from Hamilton's model, age-specific mortality curves increase exponentially across adulthood (31). This exponential increase was identified in human actuarial data by B. Gompertz in the early 19th century (60). A model bearing his name gives a fair fit to mortality data across a wide range of species (61):




Fig. 1. Female age structures modeled from life tables. Each bar shows the percentage of the population in the 5-year age class indicated in the vertical axis. Yellow bars, juvenile years; green bars, childbearing years; purple bars, post-fertile years. Humans are on the right, represented by Hadza hunter-gatherers with Blurton Jones's data (6). In this population, life expectancy at birth is 33 years. With growth rate 1.3%/year, 32% of the women (those over 15) are past the age of 45. Growing populations are younger because more are born than die. If this population was stationary, the percentage of adult women past the age of 45 would be 39% (8). The left side of the figure represents the synthetic wild chimpanzee population constructed by Hill and colleagues (63) using data from five wild study sites. Average age at first birth is 13 in wild chimpanzees so the 10- to 14-year age class is included in the childbearing years. Fertility ends by ~45 in both species. Less than 3% of the adult chimpanzees (counted as those over 10 years) are past the age of 45. The chimpanzee model assumes a stationary population.

$$m(t) = Ae^{Gt}.$$

Here m is the mortality hazard rate, G describes the rate of increase in adult mortality with increasing age (t), and A represents age-independent adult mortality. Building on previous work by G. A. Sacher (62), C. E. Finch (61) labeled A the initial mortality rate (IMR). Taking the natural log, the equation yields a line representing the logarithm of the hazard of death across adulthood with the log of the IMR as its intercept and G as its slope. In the Gompertz model, differences in longevity between populations of the same species or between species can be due to differences in the initial mortality rate (A), differences in G [or its transformed value, $\ln 2/G$, the mortality rate doubling time (MRDT)], or both. The slope (G), or the MRDT, is the demographic aging rate (62). Across species, lower initial mortality rates are correlated with shallower slopes and longer doubling times (61–64).

Some have suggested that an MRDT of 7–9 years characterizes humans (e.g., ref. 21, p. 12), but MRDTs vary at least 2-fold across human populations (28). That variation among populations is correlated with variation in the initial mortality rate. However, the correlation is in the direction opposite from that predicted by a current vs. future reproduction tradeoff. Instead of the cross-species pattern identified by Sacher (61–64), human populations with lower mortality levels (A) have faster rates of demographic aging (G). The age-specific mortality rate doubles more quickly, MRDT is shorter, when the age-independent risk of death (A) is lower.

This relationship, named for B. L. Strehler and A. S. Mildvan (26), who first identified it across human populations, is robust and well described (27). Fig. 2 shows this Strehler–Mildvan correlation across a convenience sample of human populations chosen to represent a wide range of socioecologies and initial mortality rates [from ref. 28, with two Pygmy populations (65) added here]. The figure is constructed from Gompertz models that were fitted to life tables for each population. Following Finch (61) the models consider age-specific mortality risk from ages 30 to 80 (see discussion in ref. 28). The log of A , the hazard of death at age 30 (representing the IMR) is on the horizontal axis, and G , the slope

of the log of the Gompertz curve is on the vertical axis. This correlation between the two variables across populations of the same species has also been found in widely diverse taxa where suitable data are available (27, 63). The limited data for chimpanzees are also plotted in Fig. 2. The synthetic chimpanzee population in the wild (54) used in Fig. 1 and the synthetic population from captivity (66) represent variation in IMRs and demographic aging rates in that species. The same Strehler–Mildvan relationship found across human populations holds for chimpanzees.

A Heterogeneity Hypothesis

As noted, Strehler–Mildvan correlations across populations of the same species are opposite to those generally found in cross-species comparisons. Williams' verbal arguments (10), Hamilton's formal treatment (11), and Kirkwood's disposable soma model (19) link lower mortality to stronger selection against senescence, and so slower rates of aging. Fig. 2 shows the opposite pattern. Within species lower mortality (IMR) is associated with a steeper increase in death risk across adulthood—faster rates of demographic aging. The evolutionary models all assume that more energy allocated to somatic maintenance pays off in future reproduction but leaves less for current reproductive effort. Life history variation among individuals of the same population often seems to go in the opposite direction as well. Women with higher fertility rates and later ages at last birth also have higher subsequent survival rates (23–25, 67–69). Such apparent absence of the expected tradeoffs within populations is a regular finding in field studies in animal behavior (70–72). A common explanation is that individuals differ in their resources. When these differences are ignored (or unobservable) and subjects are pooled, the resource differences obscure the tradeoff because those with more resources can have more of everything. Like houses and cars (70), more into mortgage




Fig. 2. The slope of the log of the hazard of death from age 30–80 by the log of the intercept at age 30 (IMR) taken from the values of A and G in Gompertz models calculated from life tables for a convenience sample of eleven human (yellow circles) and two synthetic chimpanzee populations (black circles). See table 1 of ref. 28 for the values plotted here. The sample includes five hunter-gatherer populations, the US and Japan to represent lower mortality levels falling in the upper left corner (the lowest IMRs and the steepest slopes), and two other cases to represent high mortality populations depending on agriculture. Here, two pygmy populations from Miglano (65) are added, the Aeta and the Batak. The chimpanzees are the synthetic wild population from Hill et al. (54) and the synthetic captive population from Dyke et al. (64). All life tables are female except for the !Kung and Agta, for which sexes were not distinguished in the original sources. Parameters were calculated on 5-year age classes, conditional on survival to the beginning of the age class preceding age 30. For the 11 human populations (yellow circles), the correlation between these estimates is -0.955 .

payments leaves less for auto loans, but those with bigger budgets can put more into both.

If there is such heterogeneity, so that health and otherwise unobserved differences in frailty vary within the populations shown in Fig. 2, that heterogeneity could account for the Strehler–Mildvan correlations in the following way (28). Frail individuals die earlier. They die even earlier under more severe conditions. Such mortality selection (73), or culling (74), changes the relative representation of subpopulations among the survivors. Older age classes are a biased subset of younger ones and that bias affects their average mortality risk. In higher mortality populations of both humans and chimpanzees, older age classes are more strongly culled leaving proportionately fewer frail survivors. Conversely, when background mortality is low, mortality selection is weaker and more of the frail survive longer. Although absolute risk of death is lower, the relative risk in each age class increases more steeply with advancing age because later age classes include more individuals with relatively greater vulnerability.

Heterogeneity could take many forms (75). One simple possibility is that populations are composed of two (unobserved) subpopulations, each with a Gompertz schedule of risk. The frailer subpopulation has higher mortality risk at each age and steeper increasing risk. The log of the risk of death at each age has both a




Fig. 3. Two model subpopulations, one frail (open green circles) and the other robust (filled blue squares), exposed to two conditions of age-independent mortality. Initial mortality rates are low (similar to the US and Japan) for the lower set of lines and high (similar to Hadza hunter-gatherers) for the upper set of lines. Both subpopulations face Gompertz age-specific risk. Initial mortality rates for the two subpopulations differ by 0.35/year in both conditions with slopes of 0.03/year and 0.085/year, respectively. The simulations plot the age-specific mortality rate for the population pooled from these two subpopulations (grey diamonds). The trendlines in red (described by the equations and correlation coefficients) measure how well the population mortality curves fit Gompertz models. The slope of the trendline is the demographic aging rate. For the population in the high mortality condition, that slope is about half as steep as in the low mortality condition. Relative size of the subpopulations at the initial adult age makes a difference. Here it is assumed to be the same at both high and low background mortalities because background risk is assumed to affect the frail proportion in two opposing ways. When age-independent mortality is high, so is the risk of early life tradeoffs that leave survivors more frail (see the discussion of early origins in the text). However, higher mortality also strengthens mortality selection across juvenile years, leaving a smaller fraction of the frail juveniles alive at maturity. On the other hand, when background mortality is low, fewer have faced early survival tradeoffs that increase frailty, making the frailer subpopulation smaller initially. Yet, weaker mortality selection across the juvenile years leaves more of the frail subpopulation surviving to adulthood.

higher intercept and higher slope in the frailer subpopulation (28, 76). Fig. 3 displays the age-specific mortality curves for simulated populations with such heterogeneity facing two different background conditions of mortality risk. In each condition there are two subpopulations, with exactly the same relative differences in age-specific risk of death. Gompertz demographic aging is linear with age on this semilog plot. The simulation uses observed ranges of variation in initial mortality rates and slopes across the sample of human populations in Fig. 2 to estimate realistic ranges. Fig. 3 shows the age-specific risk for the subpopulations and for the whole population when the subpopulations are pooled. More of the frail die at each age, and older age classes are increasingly biased toward the more robust in both conditions; but when overall mortality is low—the lower set of lines—more of the frail survive to older ages and so their higher and steeper risk has a larger effect on the relative risk of later age classes. The difference between the two subpopulations is identical in both environments, but the increase in mortality with age is about twice as steep when background mortality is lower. This is the same difference seen across empirical populations in Fig. 2.

Heterogeneity and Fertility

The same kind of differential frailty proposed to underlie the Strehler–Mildvan correlations in Fig. 2, and modeled in Fig. 3, is relevant to age-specific fertility. As shown in Fig. 1, the child-bearing years end at the age of ~45 in both humans and chimpanzees. Like other female mammals, humans and chimpanzees build initial oocyte stocks in early life that then deplete with age (77). Most of the initial stock is lost to atresia, a continuing process of cell death that begins near birth. In women, stocks decline from ~7 million oocytes at 5 months after conception to <2 million at birth and ~400,000 at puberty (78). Only one in a thousand of those remaining when ovarian cycling begins actually ovulate. Numbers continue to fall across young and middle adulthood, reaching thresholds associated first with reduced fecundability, then secondary sterility, and finally menopause ~10 years after last birth. Average ages at these thresholds differ some across populations (79) with substantial variation around the averages (80–83). The classic counts of human ovarian follicle stocks show that among females of the same age, remaining primordial follicle stocks can vary by two orders of magnitude (84–86).

Chimpanzee follicle stocks also vary among individuals of similar age (87). Archived ovarian sections taken at necropsy from captive chimpanzees of ages 0–47 years index this variation and the declining numbers with age (87). Exponential regressions fit to the age-specific primordial follicle counts on those sections and also to the whole ovary counts across that 0- to 47-year range in the classic human datasets provide a quantitative comparison of follicular loss rate in the two species. The intercepts—the heights—of the two regression lines are necessarily different because the human data represent whole ovaries and only single sections were available for the chimpanzees. [An average section is ~1/2,000 of a human ovary (84, 85)—likely the same for chimpanzees.] However, the rate of depletion with age measured this way, on these samples, across this age range, is indistinguishable between the two species (87). This similarity is consistent with a wider body of findings, including hormone and cycling data from captive chimpanzees (88–90), suggesting they would reach menopause at about the same ages humans do—if they lived long enough (91).

As implied by these similarities and noted above, humans and chimpanzees can give birth into their mid-forties but not beyond. However, in spite of this similarity in the end of the childbearing years (Fig. 1), the shapes of age-specific fertility curves in the two species are strikingly different. Fig. 4 displays the average age-specific fertilities for three hunter-gatherer populations and the conservative age-specific fertility schedule synthesized from six wild chimpanzee populations by M Emery Thompson and others (22). Human populations can differ widely in fertility levels, but

among them—hunter-gatherers included—the change in the rate of babies born to women of each age has a familiar peaked shape. “[I]n all populations where reliable records have been kept, fertility is zero until about age 15, rises smoothly to a single peak, and falls smoothly to zero by age 45–50” (ref. 92, p. 27). The fertility schedule for wild chimpanzees is flat-topped instead. The rate reached before the age of 20 continues with little change for two more decades.

The percentages running along the horizontal axis in Fig. 4 show the relative size of each age class compared to the first age class of adulthood. The chimpanzee figures come from the number of risk years observed in each age class in Emery Thompson and colleagues’ (22) supplementary table 2. For human hunter-gatherers the figures come from the female life table for Hadza foragers (6). As the percentages show, almost all of the chimpanzees that survive to adulthood then die during the childbearing years; only 1% do not. By contrast, 24% of the hunter-gatherer women die during the childbearing years; 76% do not.

Emery Thompson and colleagues (22) demonstrated heterogeneity in chimpanzee fertility in their six-site sample by looking for associations between fertility rates and survival in females over the age of 25. They divided their observations into healthy and unhealthy years. An observation year for a given chimpanzee was considered healthy if she survived an additional 5 years or more, and unhealthy if she did not. Their figure 2 (ref. 22, p. 2152) shows that fertility in the thirties was about twice as high in females who would survive at least 5 more years than in those who would not. The finding indicates that mortality selection across the childbearing years culls the females with lower fertility. As the age classes shrink to almost nothing, they are increasingly biased to the less frail, more fertile females. Consequently, average fertility changes little even if the fertility of the survivors is declining relative to their own earlier rate.

We found similar heterogeneity in fertility in 19th century Utah women [the Utah Population Data Base (UPDB) (93)]. Although not hunter-gatherers, these women practiced natural fertility (94),




Fig. 4. Age-specific fertility rates (ASFR) for humans and chimpanzees. Humans (red circles) are represented by the average of three hunter-gatherer populations: !Kung Bushmen of Botswana (4), Ache of Paraguay (4), and Hadza of Tanzania (6). Estimates for chimpanzees in the wild (blue squares) come from the conservative fertility schedule synthesized from six study sites by Emery Thompson et al. (22). The bumps reflect small sample size (627 risk years in the initial chimpanzee adult age class declining to 8 risk years in the 45- to 49-year interval (ref. 22, supplementary table 2). The percentages along the horizontal axis indicate the proportion of those reaching adulthood that survive to the age class. The top row of percentages are estimates for chimpanzees from the number of risk years in each age class (ref. 22, supplementary table 2). They are just slightly lower than the model in Fig. 1 from the life table (54). The bottom row is human estimates from the female life table for Hadza hunter-gatherers (6).

so potential for continued child bearing is reflected by actual births. Individual records make it possible to investigate links between variation in fertility rate and age at last birth. Of 42,493 parous UPDB women born between 1849 and 1890, the 10,440 whose fertility ended before the age of 35 had fertility rates in the preceding years about half as high as the 2,695 women who would have last births after 45 (95). This parallels the chimpanzee variation with an important difference: all the women in the Utah sample, whatever their age of last birth, survived at least to the age of 50. The sample was restricted to women who lived at least to that age to avoid the confound of early last births due to early death (96). Subjects were also restricted to those married once and neither widowed nor divorced to reduce effects these characteristics may have on fertility.

Assuming that heterogeneity in fertility is similar in the hunter-gatherer women, this variation combined with the different survival schedules of humans and chimpanzees can explain the different shapes of the fertility schedules shown in Fig. 4. Most women, whatever their frailty, survive the childbearing years, whereas across those years mortality culls chimpanzee females down to a least frail few. The human schedule is peaked because women with both high and low risk of fertility failure outlive the childbearing years. Beginning about the age of 30, each subsequent age interval contains more women who are past their last parturition. This drives down the average rate of baby production for later age intervals (95, 97, 98). The chimpanzee schedule is flat because heterogeneity in ovarian aging is culled away by mortality selection. Most chimpanzees die during the childbearing years and the survivors are females whose fertility rate has been high all along (22, 95). In captive chimpanzees, lower mortality allows more frail individuals to survive longer so that captive chimpanzee fertility slopes down from a peak, more like the human pattern (99, 100).

Origins of Heterogeneity in Early Life

Human age structures looked much like the hunter-gatherer example shown in Fig. 1 until the 20th century when life expectancies at birth began to increase in some populations (1, 2). Until the mid-20th century, these increases were largely a consequence of decreasing numbers of dying infants and children: lower juvenile mortality is strongly associated with lower fertility. Fig. 5 shows number of births for UPDB women who survived at least to 50 by their own birth year across the 19th century (96). After the middle of the 19th century, fertility began a steady decline—falling to half of its earlier level by 1900. Fig. 5 also shows a concurrent change in adult mortality. The average age at death for women who had survived at least to 50 increased from ~75 at mid-century to ~80 at its end. These decreases in mortality and fertility typify changes in some other populations at about the same time, likely due to improvements in nutrition, sanitation, and medicine (21, 101). By the end of the 20th century, continuing decreases in fertility and increases in juvenile and adult survival resulted in life expectancies double those of most historical and ethnographic populations (1, 9, 102). The increases in survival allowed increased heterogeneity at older ages. Other effects on heterogeneity are likely as well.

Associations between regional infant mortality rates and late life morbidities led D. J. Barker to propose his infant and fetal origins of adult disease hypothesis (103–105). Those who survive nutritional and disease insults in early life are predisposed to metabolic and cardiovascular disease in adulthood. Pursuit of Barker’s hypothesis has revealed that when historical cohorts have been exposed to famines and epidemics during fetal life they have higher rates of disease and mortality in later adulthood than do adjacent cohorts (106, 107). Analyses of adult morbidity and mortality by birth month and season have yielded similar evidence of heterogeneity in frailty stemming from nutritional constraints and disease exposure in early life (108, 109).

Differences between cohorts are necessarily an underestimate of the likely heterogeneity. B. Mamzunder and colleagues (107) compared morbidities in later adulthood between Americans who




Fig. 5. Number of births (green circles) and age at death (diamonds) in cohorts of UPDB women by their birth year across the 19th century (ref. 150, redrawn from ref. 96). Only women who survived past the age of 50 are included.

were likely exposed in fetal life to the 1918 influenza pandemic and those in adjacent cohorts. As they noted, “Maternal health during the pandemic peak... varied widely from no clinical infection, mild uncomplicated flu or flu with severe secondary pneumonia that still permitted normal birth” (p. 4). Those whose birth dates indicate probable fetal flu exposure must include some unexposed individuals. In the same way, adjacent cohorts must include some individuals whose mothers experienced infection. Nutrition, energy expenditure, and stress may also impact the effects of disease (21, 110, 111), and recent disease history of groups in the sample may also influence responses to early life conditions (112, 113). Even with the imperfect association between exposure and birth date and effects of these unmeasured covariates, rates of cardiovascular disease after the age of 60 were >20% higher in those whose fetal development coincided with pandemic (107). That this is a minimum estimate of early life effects on heterogeneity in aging rates is underscored by longitudinal datasets documenting within-cohort associations between early growth and both ovarian aging and mid-life physical performance (114, 115).

The early origins hypothesis predicts that declines in mortality in the Utah women during the second half of the 19th century would affect the next generation. Longer survival likely indicates better nourishment, less illness, and reduced hardship. If so, the children of those surviving longer would have been less exposed to nutritional limits and infection in early life, and so have lower risks of various later morbidities. Improvements in nutrition, general public health and subsequently medical interventions should mitigate early life insults and reduce consequent heterogeneity. However, lowered mortality also reduces mortality selection, allowing greater heterogeneity to persist to older ages. This heterogeneity hypothesis (Fig. 3) to explain population variation in demographic aging (Fig. 2) applies to chimpanzees (and other taxa) as well as humans. Life expectancy at birth for female chimpanzees in the wild is 15 years (54). In captivity it is 29 years (66). This doubling of chimpanzee life expectancy is associated with reductions in rates of infection and nutritional stress (38, 116). In both chimpanzees and humans, improvements in nutrition and hygiene combined with medical interventions can double life expectancy. And in both, longer life expectancies are associated with faster rates of demographic aging (Fig. 2)—due perhaps, as argued here, to increased heterogeneity of frailty at older ages (Fig. 3).

Back to Grandmothers

This heterogeneity hypothesis may explain why humans, chimpanzees, and other taxa display Strehler–Mildvan correlations. The similarities cannot explain why humans usually outlive the childbearing years and chimpanzees do not (Figs. 1 and 2). Physiological mechanisms, let alone genetic differences that underlie the

survival differences, remain elusive (102, 117), although mitochondrial mutation rates may be involved (118, 119). Hamilton’s forces (11, 33) do not specify particular mechanisms of aging, but their incorporation in an analysis of human survival curves points to a deep history of reproductive benefits accruing to postmenopausal women in our lineage. Mueller, Rose, and Rauser have focused attention on the period of life in many species when mortality rates slow from an exponential increase and may become constant at succeeding age intervals. They reject Vaupel’s heterogeneity of frailty hypothesis (30) as a general explanation for these mortality plateaus, finding evidence more consistent with expectations from evolutionary theory about late life (31–33). When individuals survive past normal lifespans, they are beyond the ages where senescence has been molded by ancestral forces of selection. “Hamiltonian theory predicts that late-life mortality rates should plateau and evolve according to the last age of reproduction in a population’s evolutionary history” (ref. 32, p. 26). Because human mortality rates begin to decelerate and depart from a Gompertz curve only around the ninetieth year (30), the mortality plateau criterion implies that contributions to reproduction from ancestral grandmothers continued through their eighties.

This demographic evidence of grandmaternal effects on reproduction in our lineage has other implications that can barely be touched on here. S. B. Hrdy (120–123) has hypothesized that selection pressures for distinctively human cognitive and emotional capacities arose from our evolution as cooperatively breeding apes. Unlike our nearest living relatives, human mothers accept help with babies right from parturition. Depending on help, they can bear a new baby while previous offspring still need provisioning. This has consequences for selection pressures on both mothers and infants. Unlike chimpanzee mothers, humans must also consider the occupation and whereabouts of potential helpers as well as the needs of still dependent weaned children. Abilities to juggle these additional concerns supersede the more single-minded focus on the newborn of other ape mothers. The novel maternal sensitivities create problems in turn for human infants that do not arise for other infant apes. Human babies cannot count on mother’s undivided commitment, so capacities to actively engage her and also to evaluate and engage other helpers are crucial. In high infant mortality environments selection on those capacities would have been especially strong. Hrdy (122) links those circumstance to the evolution in our lineage of motivations and capacities for intersubjective engagement that M. Tomasello and colleagues (124, 125) identify as the foundation for human prosociality.

Ethnographers have documented the ubiquity and importance of allomothering from many kinds of kin in living human communities (126), but grandmothers in particular are implicated in the hypothesis about the evolution of human life history entertained here. If ancestral grandmothers provided the help that initially allowed mothers in our lineage to move on to the next baby before the previous one could feed itself, propelling the evolution of human postmenopausal longevity, that initiated cooperative breeding in a previously independently breeding ancestral ape. These arguments link distinctively human cognitive and emotional capacities to selection pressures that arose as a consequence of ancestral grandmothering.

Ovarian aging appears to differ little between modern humans and chimpanzees, making it likely the same pattern characterized our ancestors. Before the shifts to greater longevity in our lineage, heterogeneity in ovarian and somatic aging would have been strongly culled by mortality selection across the childbearing years. If grandmother effects reduced mortality across those years, heterogeneity in ovarian aging would have expanded as more and more women outlived their fertility. Subsidies for relatives’ reproduction would have begun well before the average age at last birth, let alone the average age of menopause. By this argument, heterogeneity in ovarian aging is an ancient legacy of grandmothering in our lineage; but now such heterogeneity poses

unprecedented concerns in the human populations where child-bearing is delayed and nuclear families are isolated as never before. Many women find they have missed their own windows of fertility (83). Although aging is often seen as a process that befalls the old, evolutionary theories of aging predict that function begins to decline in early adulthood. Such declines have been documented not only in fertility, but in muscle strength and cognitive performance (127, 128); and where mortality levels have dropped to evolutionarily unprecedented lows, heterogeneity in somatic competence is increasingly well documented in those past mid-life (129, 130). Just as grandmothing may have expanded heterogeneity in ovarian aging by lowering mortality across the child-

bearing years, recently dropping mortality rates at older ages expand heterogeneity well beyond them. As continuing innovations in medical and daily living technologies interact with mortality selection to produce complex dynamics in the health and welfare of elders (131), the heterogeneity in ovarian and somatic aging that is an aspect of our evolved life history becomes an increasing medical as well as social, economic, and political concern of our time.

ACKNOWLEDGMENTS. I am grateful for valuable insights and advice from J. F. O'Connell, J. E. Coxworth, J. K. Blevins, C. W. Kuzawa, and S. B. Hrdy, and for research support from the National Science Foundation.

- Oeppen J, Vaupel JW (2002) Demography. Broken limits to life expectancy. *Science* 296:1029–1031.
- Keyfitz N, Fleiger W (1968) *World Population: An Analysis of Vital Data* (Univ of Chicago Press, Chicago).
- Howell N (1979) *Demography of the Dobe! Kung* (Academic Press, New York).
- Hill K, Hurtado AM (1996) *Ache Life History: The Ecology and Demography of a Foraging People* (de Gruyter, New York).
- Early JD, Headland TN (1998) *Population Dynamics of a Philippine Rain Forest People: The San Ildefonso Agta* (Univ Press of Florida, Gainesville).
- Blurton Jones NG, Hawkes K, O'Connell JF (2002) Antiquity of postreproductive life: Are there modern impacts on hunter-gatherer postreproductive life spans? *Am J Hum Biol* 14:184–205.
- Hill K, Hurtado AM, Walker RS (2007) High adult mortality among Hiwi hunter-gatherers: Implications for human evolution. *J Hum Evol* 52:443–454.
- Hawkes K, Blurton Jones NG (2005) Human age structures, paleodemography, and the grandmother hypothesis. *Grandmotherhood: The Evolutionary Significance of the Second Half of Female Life*, eds Roth G, Voland E, Chasiotis A, Schiefenhovel W (Rutgers Univ Press, New Brunswick, NJ), pp 118–140.
- Gurven M, Kaplan H (2007) Longevity among hunter-gatherers: A cross-cultural examination. *Popul Dev Rev* 33:321–365.
- Williams GC (1957) Pleiotropy, natural selection, and the evolution of senescence. *Evolution* 11:398–411.
- Hamilton WD (1966) The moulding of senescence by natural selection. *J Theor Biol* 12:12–45.
- Hawkes K, O'Connell JF, Jones NG, Alvarez H, Charnov EL (1998) Grandmothering, menopause, and the evolution of human life histories. *Proc Natl Acad Sci USA* 95:1336–1339.
- Robson SL, van Schaik CP, Hawkes K (2006) The derived features of human life history. *The Evolution of Human Life History*, eds Hawkes K, Paine RR (School of American Research Press, Santa Fe, NM), pp 17–45.
- Charnov EL (1991) Evolution of life history variation among female mammals. *Proc Natl Acad Sci USA* 88:1134–1137.
- Charnov EL (1993) *Life History Invariants: Some Explorations of Symmetry in Evolutionary Ecology* (Oxford Univ Press, Oxford).
- Williams GC (1966) Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am Nat* 100:687–690.
- Stearns SC (1992) *The Evolution of Life Histories* (Oxford Univ Press, Oxford).
- Charlesworth B (1994) *Evolution in Age-structured Populations* (Cambridge Univ Press, Cambridge, U.K.).
- Kirkwood TBL, Rose MR (1991) Evolution of senescence: Late survival sacrificed for reproduction. *Proc R Soc Lond B Biol Sci* 332:15–24.
- Kirkwood TBL, Holliday R (1979) The evolution of ageing and longevity. *Proc R Soc Lond B Biol Sci* 205:532–546.
- Finch CE (2007) *The Biology of Human Longevity: Inflammation, Nutrition, and Aging in the Evolution of Lifespans* (Academic Press, Amsterdam).
- Emery Thompson M, et al. (2007) Aging and fertility patterns in wild chimpanzees provide insights into the evolution of menopause. *Curr Biol* 17:2150–2156.
- Perls TT, Alpert L, Fretts RC (1997) Middle-aged mothers live longer. *Nature* 389:133.
- Jacobsen BK, Heuch I, Kvåle G (2003) Age at natural menopause and all-cause mortality: A 37-year follow-up of 19,731 Norwegian women. *Am J Epidemiol* 157:923–929.
- Gagnon A, et al. (2009) Is there a trade-off between fertility and longevity? A comparative study of women from three large historical databases accounting for mortality selection. *Am J Hum Biol* 21:533–540.
- Strehler BL, Mildvan AS (1960) General theory of mortality and aging. *Science* 132:14–21.
- Gavrilov LA, Gavrilova NS (2001) The reliability theory of aging and longevity. *J Theor Biol* 213:527–545.
- Hawkes K, Smith KR, Robson SL (2009) Mortality and fertility rates in humans and chimpanzees: How within-species variation complicates cross-species comparisons. *Am J Hum Biol* 21:578–586.
- Vaupel JW, Manton KG, Stallard E (1979) The impact of heterogeneity in individual frailty on the dynamics of mortality. *Demography* 16:439–454.
- Vaupel JW, et al. (1998) Biodemographic trajectories of longevity. *Science* 280:855–860.
- Mueller LD, Rose MR (1996) Evolutionary theory predicts late-life mortality plateaus. *Proc Natl Acad Sci USA* 93:15249–15253.
- Rausser CL, Mueller LD, Rose MR (2006) The evolution of late life. *Ageing Res Rev* 5:14–32.
- Rose MR, Rausser CL, Benford G, Matos M, Mueller LD (2007) Hamilton's forces of natural selection after forty years. *Evolution* 61:1265–1276.
- Wood B (2010) Reconstructing human evolution: Achievements, challenges, and opportunities. *Proc Natl Acad Sci* 107:8902–8909.
- Glazko GV, Nei M (2003) Estimation of divergence times for major lineages of primate species. *Mol Biol Evol* 20:424–434.
- Wood BA, Collard M (1999) The human genus. *Science* 284:65–71.
- Robson SL, Wood BA (2008) Hominin life history: Reconstruction and evolution. *J Anat* 212:394–425.
- Goodall J (1986) *The Chimpanzees of Gombe* (Harvard Univ Press, Cambridge, MA).
- Blurton Jones NG, Hawkes K, O'Connell JF (1997) Why do Hadza children forage? *Uniting Psychology and Biology: Integrative Perspectives on Human Development*, eds Segal N, Weisfeld GE, Weisfeld CC (Am Psychol Assoc, Washington, DC), pp 279–313.
- Bliege Bird R, Bird DW (2002) Constraints of knowing or constraints of growing? Fishing and collecting among the children of Mer. *Hum Nat* 13:297–321.
- Bird DW, Bliege-Bird R (2005) Martu children's hunting strategies in the Western Desert, Australia: Foraging and the evolution of human life histories. *Hunter Gatherer Childhoods*, eds Hewlett BS, Lamb ME (de Gruyter, New York), pp 129–146.
- Hawkes K, O'Connell JF, Blurton Jones NG (1995) Hadza children's foraging: Juvenile dependency, social arrangements and mobility among hunter-gatherers. *Curr Anthropol* 36:688–700.
- Hawkes K, O'Connell JF, Blurton Jones NG (1997) Hadza women's time allocation, offspring provisioning and the evolution of post-menopausal lifespans. *Curr Anthropol* 38:551–577.
- Hawkes K, O'Connell JF, Blurton Jones NG (2003) Human life histories: Primate tradeoffs, grandmothering socioecology, and the fossil record. *Primate Life Histories & Socioecology*, eds Kappeler P, Pereira M (Univ of Chicago Press, Chicago), pp 204–227.
- Hawkes K (2003) Grandmothers and the evolution of human longevity. *Am J Hum Biol* 15:380–400.
- O'Connell JF, Hawkes K, Blurton Jones NG (1999) Grandmothering and the evolution of Homo erectus. *J Hum Evol* 36:461–485.
- deMenocal PB (1995) Plio-Pleistocene African climate. *Science* 270:53–59.
- Bromage TG, Schrenk F (1999) *African Biogeography, Climate Change, and Human Evolution* (Oxford Univ Press, Oxford).
- O'Connell JF, Hawkes K, Lupo KD, Blurton Jones NG (2002) Male strategies and Plio-Pleistocene archaeology. *J Hum Evol* 43:831–872.
- Kaplan HS, Robson AJ (2009) We age because we grow. *Proc R Soc Lond B Biol Sci* 276:1837–1844.
- Lee RD (2003) Rethinking the evolutionary theory of aging: Transfers, not births, shape senescence in social species. *Proc Natl Acad Sci USA* 100:9637–9642.
- Lee RD (2008) Sociality, selection, and survival: Simulated evolution of mortality with intergenerational transfers and food sharing. *Proc Natl Acad Sci USA* 105:7124–7128.
- Alvarez HP (2000) Grandmother hypothesis and primate life histories. *Am J Phys Anthropol* 133:435–450.
- Hill K, et al. (2001) Mortality rates among wild chimpanzees. *J Hum Evol* 39:1–14.
- Sahlins MD (1959) The social life of monkeys, apes and primitive man. *Hum Biol* 31:54–73.
- Kaplan HK (1994) Evolutionary and wealth flows theories of fertility: Empirical tests and new models. *Popul Dev Rev* 20:753–791.
- Blurton Jones NG, Marlowe F (2002) Selection for delayed maturity: Does it take 20 years to learn to hunt and gather? *Hum Nat* 13:199–238.
- Walker R, Hill K (2003) Modeling growth and senescence in physical performance among the Ache of eastern Paraguay. *Am J Hum Biol* 15:196–208.
- Finch CE, Stanford CB (2004) Meat-adaptive genes and the evolution of slower aging in humans. *Q Rev Biol* 79:3–50.
- Gompertz B (1825) On the nature of the function expressive of the law of human mortality. *Proc R Soc Lond B Biol Sci* 115:513–585.
- Finch CE (1990) *Longevity, Senescence, and the Genome* (University of Chicago Press, Chicago).
- Sacher GA (1977) Life table modification and life prolongation. *Handbook of the Biology of Aging*, eds Finch CE, Hayflick L (Van Nostrand, New York), pp 282–263.
- Pletcher SD, Neuhäuser C (2000) Biological aging—criteria for modeling and a new mechanistic model. *Int J Mod Phys C* 11:525–546.

64. Ricklefs RE (1998) Evolutionary theories of aging: Confirmation of a fundamental prediction, with implications for the genetic basis and evolution of life span. *Am Nat* 152:24–44.
65. Migliano AB (2005) *Why are Pygmies Small: Ontogenetic Implications of Life History Evolution*. PhD dissertation (Cambridge Univ, Cambridge, U.K.).
66. Dyke B, Gage TB, Alford PL, Swenson B, Williams-Blangero S (1995) Model life table for captive chimpanzees. *Am J Primatol* 37:25–37.
67. Smith KR, Mineau GP, Bean LL (2002) Fertility and post-reproductive longevity. *Soc Biol* 49:185–205.
68. Müller H-G, Chiou J-M, Carey JR, Wang J-L (2002) Fertility and life span: Late children enhance female longevity. *J Gerontol A Biol Sci Med Sci* 57:B202–B206.
69. Smith KR, et al. (2009) Familial aggregation of survival and late female reproduction. *J Gerontol A Biol Sci Med Sci* 64:740–744.
70. van Noordwijk AJ, de Jong G (1986) Acquisition and allocation of resources: Their influence on life history traits. *Am Nat* 128:137–142.
71. Pettifor RA, Perrins CM, McCleery RH (1988) Individual optimization of clutch size in Great Tits. *Nature* 336:160–162.
72. Lessels CM (1991) The evolution of life histories. *Behavioural Ecology: An Evolutionary Approach*, eds Krebs JR, Davies NB (Blackwell Scientific, Oxford), pp 32–68.
73. Manton KG, Stallard E (1984) *Recent Trends in Mortality Analysis* (Academic Press, Orlando).
74. Wachter KW (2003) Hazard curves and life span prospects. *Popul Dev Rev* 29 (3 Suppl):270–291.
75. Vaupel JW, Yashin AI (1985) Heterogeneity's ruses: Some surprising effects of selection on population dynamics. *Am Stat* 39:176–185.
76. Wilmoth JR, Horiuchi S (1999) Do the oldest old grow old more slowly? *The Paradoxes of Longevity*, eds Forette B, Franceschi C, Robine JM, Allard M (Springer, Heidelberg), pp 25–60.
77. vom Saal FS, Finch CE, Nelson JF (1994) Natural history and mechanisms of reproductive aging in humans, laboratory rodents, and other selected vertebrates. *The Physiology of Reproduction*, eds Knobil E, Neill JD (Raven, New York), 2nd Ed, pp 1213–1314.
78. Baker TG (1963) A quantitative and cytological study of germ cells in human ovaries. *Proc R Soc Lond B Biol Sci* 158:417–433.
79. Bentley GR, Muttukrishna S (2007) Potential use of biomarkers for analyzing interpopulation and cross-cultural variability in reproductive aging. *Menopause* 14: 668–679.
80. Faddy MJ, Gosden RG (1996) A model conforming the decline in follicle numbers to the age of menopause in women. *Hum Reprod* 11:1484–1486.
81. O'Connor KA, Holman DJ, Wood JW (2001) Menstrual cycle variability and the perimenopause. *Am J Hum Biol* 13:465–478.
82. Sievert LL (2006) *Menopause: A Biocultural Perspective* (Rutgers Univ Press, New Brunswick, NJ).
83. Broekmans FJ, Soules MR, Fauser BC (2009) Ovarian aging: Mechanisms and clinical consequences. *Endocr Rev* 30:465–493.
84. Block E (1952) Quantitative morphological investigations of the follicular system in women; variations at different ages. *Acta Anat (Basel)* 14:108–123.
85. Richardson SJ, Senikas V, Nelson JF (1987) Follicular depletion during the menopausal transition: Evidence for accelerated loss and ultimate exhaustion. *J Clin Endocrinol Metab* 65:1231–1237.
86. Gougeon A, Ecochard R, Thalabard JC (1994) Age-related changes of the population of human ovarian follicles: Increase in the disappearance rate of non-growing and early-growing follicles in aging women. *Biol Reprod* 50:653–663.
87. Jones KP, et al. (2007) Depletion of ovarian follicles with age in chimpanzees: Similarities to humans. *Biol Reprod* 77:247–251.
88. Graham CE (1979) Reproductive function in aged female chimpanzees. *Am J Phys Anthropol* 50:291–300.
89. Gould KG, Flint M, Graham CE (1981) Chimpanzee reproductive senescence: A possible model for evolution of the menopause. *Maturitas* 3:157–166.
90. Lacreuse A, et al. (2008) Menstrual cycles continue into advanced old age in the common chimpanzee (*Pan troglodytes*). *Biol Reprod* 79:407–412.
91. Walker ML, Herndon JG (2008) Menopause in nonhuman primates? *Biol Reprod* 79: 398–406.
92. Coale AJ, Demeny P (1983) *Regional Model Life Tables and Stable Populations* (Academic Press, New York), 2nd Ed.
93. Bean LL, Mineau GP, Anderton DL (1990) *Fertility Change on the American Frontier, Adaptation and Innovation* (Univ of California Press, Berkeley).
94. Henry L (1961) Some data on natural fertility. *Eugen Q* 18:81–91.
95. Hawkes K, Smith KR (2010) Do women stop early? Similarities in fertility decline between humans and chimpanzees. *Ann NY Acad Sci*, 10.1111/j.1749-6632.2010.05527.x.
96. Hawkes K, Smith KR (2009) Brief communication: Evaluating grandmother effects. *Am J Phys Anthropol* 140:173–176.
97. Wood JW (1989) Fecundity and natural fertility in humans. *Oxf Rev Reprod Biol* 11: 61–109.
98. Wood JW (1994) *Dynamics of Human Reproduction: Biology, Biometry, Demography* (de Gruyter, New York).
99. Roof KA, Hopkins WD, Izard MK, Hook M, Schapiro SJ (2005) Maternal age, parity, and reproductive outcome in captive chimpanzees (*Pan troglodytes*). *Am J Primatol* 67:199–207.
100. Littleton J (2005) Fifty years of chimpanzee demography at Taronga Park Zoo. *Am J Primatol* 67:281–298.
101. Fogel R (2004) *The Escape from Hunger and Premature Death 1700–2100: Europe, America and the Third World* (Cambridge Univ Press, Cambridge, U.K.).
102. Finch CE (2010) Evolution of the human lifespan and diseases of aging: Roles of infection, inflammation, and nutrition. *Proc Natl Acad Sci USA* 107 (Suppl 1): 1718–1724.
103. Barker DJ, Osmond C (1986) Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet* 1:1077–1081.
104. Barker DJ, Osmond C, Golding J, Kuh D, Wadsworth ME (1989) Growth in utero, blood pressure in childhood and adult life, and mortality from cardiovascular disease. *BMJ* 298:564–567.
105. Gluckman PD, Hanson MA, Cooper C, Thornburg KL (2008) Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med* 359:61–73.
106. Finch CE, Crimmins EM (2004) Inflammatory exposure and historical changes in human life-spans. *Science* 305:1736–1739.
107. Mazumder B, Almond S, Park K, Crimmins EM, Finch CE (2009) Lingering prenatal effects of the 1918 influenza pandemic on cardiovascular disease. *J Dev Origin Health Dis* 1:1–9.
108. Moore SE, et al. (1997) Season of birth predicts mortality in rural Gambia. *Nature* 388:434.
109. Doblhammer G, Vaupel JW (2001) Lifespan depends on month of birth. *Proc Natl Acad Sci USA* 98:2934–2939.
110. Kuzawa CW, Quinn EA (2009) Developmental origins of adult function and health: Evolutionary hypotheses. *Annu Rev Anthropol* 38:131–147.
111. Kuzawa CW, Sweet CW, et al. (2009) Epigenetics and the Embodiment of Race: Developmental Origins of US Racial Disparities in Cardiovascular Health. *Am J Hum Biol* 21:2–16.
112. Costa DL, Helmchen LA, Wilson S (2007) Race, infection, and arteriosclerosis in the past. *Proc Natl Acad Sci USA* 104:13219–13224.
113. Pennington R, Gatenbee C, Kennedy B, Harpending H, Cochran G (2009) Group differences in proneness to inflammation infection. *Genet Evol* 9:1371–1380.
114. Hardy R, Kuh D (2002) Does early growth influence timing of the menopause? Evidence from a British birth cohort. *Hum Reprod* 17:2474–2479.
115. Kuh D, et al. (2002) Birth weight, childhood size, and muscle strength in adult life: Evidence from a birth cohort study. *Am J Epidemiol* 156:627–633.
116. Williams JM, et al. (2008) Causes of death in the Kasekela chimpanzees of Gombe national Park, Tanzania. *Am J Primatol* 70:766–777.
117. de Magalhães JP, Church GM (2007) Analyses of human-chimpanzee orthologous gene pairs to explore evolutionary hypotheses of aging. *Mech Ageing Dev* 128: 355–364.
118. Kujoth GC, Bradshaw PC, Haroon S, Prolla TA (2007) The role of mitochondrial DNA mutations in mammalian aging. *PLoS Genetics* 3 (3):e24.
119. Nabholz B, Glemin S, Galtier N (2007) Strong variations of mitochondrial mutation rate across mammals—the longevity hypothesis. *Mol Biol Evol* 25:120–130.
120. Hrdy SB (1999) *Mother Nature: A History of Mothers, Infants, and Natural Selection* (Pantheon, New York).
121. Hrdy SB (2005) Comes the child before the man: How cooperative breeding and prolonged postweaning dependence shaped human potentials. *Hunter Gatherer Childhoods*, eds Hewlett B, Lamb M (Transaction, New Brunswick, NJ), pp 65–91.
122. Hrdy SB (2009) *Mothers and Others: The Evolutionary Origins of Mutual Understanding* (Harvard Univ Press, Cambridge, MA).
123. Burkhardt JM, Hrdy SB, van Schaik CP (2009) Cooperative Breeding and human cognitive evolution. *Evol Anthropol* 18:175–186.
124. Tomasello M, Carpenter M, Call J, Behne T, Moll H (2005) Understanding and sharing intentions: The origins of cultural cognition. *Behav Brain Sci* 28:675–691, discussion 691–735.
125. Tomasello M, Rakoczy H (2003) What makes human cognition unique? From individual to shared to collective intentionality. *Mind & Language* 18:121–147.
126. Sear R, Mace R (2008) Who keeps children alive? A review of the effects of kin on child survival. *Evol Hum Behav* 29:1–18.
127. Hunter SK, Thompson MW, Adams RD (2000) Relationships among age-associated strength changes and physical activity level, limb dominance, and muscle group in women. *J Gerontol A Biol Sci Med Sci* 55A:B264–B273.
128. Salthouse TA (2009) When does age-related cognitive decline begin? *Neurobiol Aging* 30:507–514.
129. Mitnitski A, et al. (2005) Relative Fitness and Frailty of Elderly Men and Women in Developed Countries and Their Relationship with Mortality. *J Am Gerontol Soc* 53: 2184–2189.
130. Rockwood K, Mitnitski A (2007) Frailty in Relation to the Accumulation of Deficits. *J Gerontol Med Sci* 62A:722–727.
131. Manton KG (2008) Recent declines in chronic disability in the elderly U.S. population: Risk factors and future dynamics. *Annu Rev Public Health* 29:91–113.